

JEDNAČINE MATEMATIČKE FIZIKE - januar 2002

1. Rešiti Košijev problem u  $I$  kvadrantu

$$\begin{cases} x^2 u_{xx} - y^2 u_{yy} - 2yu_y = 0 \\ u(1, y) = y \\ u_x(1, y) = y \end{cases}$$

2. Rešiti mešoviti problem

$$\begin{cases} u_{tt} = u_{xx} - 4u, \quad 0 < x < 1, \quad t > 0 \\ u(0, t) = 0 \\ u_x(1, t) = 0 \\ u(x, 0) = x(x - 1) \\ u_t(x, 0) = 0 \end{cases}$$

3. Rešiti mešoviti problem

$$\begin{cases} u_t = u_{xx} + u - \left(x - \frac{\pi}{2}\right)t^2 + t(1 - \pi + 2x) - 1 + 2\sin 2x \sin x, \quad 0 < x < \frac{\pi}{2}, \quad t > 0 \\ u_x(0, t) = t^2 \\ u\left(\frac{\pi}{2}, t\right) = t \\ u(x, 0) = 2\cos x \end{cases}$$

JEDNAČINE MATEMATIČKE FIZIKE - januar 2002

1. Rešiti Košijev problem u  $I$  kvadrantu

$$\begin{cases} x^2 u_{xx} - y^2 u_{yy} - 2yu_y = 0 \\ u(1, y) = y \\ u_x(1, y) = y \end{cases}$$

2. Rešiti mešoviti problem

$$\begin{cases} u_{tt} = u_{xx} - 4u, \quad 0 < x < 1, \quad t > 0 \\ u(0, t) = 0 \\ u_x(1, t) = 0 \\ u(x, 0) = x(x - 1) \\ u_t(x, 0) = 0 \end{cases}$$

3. Rešiti mešoviti problem

$$\begin{cases} u_t = u_{xx} + u - \left(x - \frac{\pi}{2}\right)t^2 + t(1 - \pi + 2x) - 1 + 2\sin 2x \sin x, \quad 0 < x < \frac{\pi}{2}, \quad t > 0 \\ u_x(0, t) = t^2 \\ u\left(\frac{\pi}{2}, t\right) = t \\ u(x, 0) = 2\cos x \end{cases}$$